

Why do we need supercomputers to understand the electrocardiographic T wave ?

Mark Potse, A.-Robert LeBlanc, Alain Vinet

Research Center, Hôpital du Sacré-Coeur, Montréal,

Department of Physiology, Institute of Biomedical Engineering, Université de Montréal, Québec, Canada

ABSTRACT

Objective: Propagation of depolarisation and repolarisation in myocardium results from an interplay of membrane potential, transmembrane current, and intercellular current. This process can be represented mathematically with a reaction-diffusion (RD) equation. Solving RD equations for a whole heart requires a supercomputer. Therefore, earlier models used predefined action potential (AP) shapes and fixed propagation velocities. We discuss why RD models are important when T waves are studied.

Methods: We simulated propagating AP with an RD model of the human heart, which included heterogeneity of membrane properties. Computed activation times served as input to a model that used predefined AP, and to a "hybrid model" that computed AP only during repolarisation. The hybrid model was tested with different spatial resolutions. Electrocardiograms (ECGs) were computed with all three models.

Results: Computed QRS complexes were practically identical in all models. T waves in the fixed-AP model had 20 to 40% larger amplitudes in leads V1–V3. The hybrid model produced the same T waves as the RD model at 0.25-mm resolution, but underestimated T-wave amplitude at lower resolutions.

Conclusion: Fixed AP waveforms in a forward ECG model lead to exaggerated T waves. Hybrid models require the same high spatial resolution as RD models. (*Anadolu Kardiyol Derg 2007; 7 Suppl 1; 123-4*)

Key words: ventricular repolarisation, computer model

Introduction

Mathematical models have played a prominent role in the study of ion channels in the cardiac cell membrane, where a succession of models, experiments, and refutations has led to the theories that are currently successful in explaining the action potential (AP) (1). Accurate models for human ventricular myocytes are now available (2, 3). By connecting such models together one can create a model that simulates propagating action potentials in the heart. Such models are called reaction-diffusion (RD) models, where "reaction" represents the ionic current sources and "diffusion" is the mathematical description of the current spread through the tissue.

Because of the steepness of the myocardial AP upstroke, and the consequent steepness of the spatial distribution of transmembrane potential, RD models must have a high spatial resolution of 0.1 mm to 0.25 mm. Thus, modeling an entire (human) heart with such models requires evaluation of potentials at 12 million points or more (4). Such models require at least 8 GB memory - twice as much as a personal computer can possibly have - and take several days of processor time to compute a single heart beat. Supercomputers with tens or hundreds of processors and very large memory are typically used to run such models (5).

A more economical way to model the entire heart is to assume a fixed AP shape and fixed propagation velocities (6). This removes the need to evaluate ionic currents and diffusion currents at millions of sites, and allows the use of a much lower spatial resolution (in the order of 1 mm). The disadvantage of this approach is that AP changes

due to electrical current flow between cells with different characteristics or different activation times are not taken into account.

One of the compromises that can be made between reaction-diffusion and fixed AP models is a RD model at low spatial resolution. Because such a model would not be able to simulate propagated depolarization, its elements must be stimulated at fixed activation times. The advantage over a fixed-AP model is that the effect of cellular coupling is taken into account during the plateau and repolarisation phase of the AP.

We compared the accuracy of such models for the purpose of electrocardiogram (ECG) simulation.

Methods

We simulated propagating AP with an RD model of the human heart at 0.25-mm resolution. Details of this model have been described previously (4, 5). Ionic currents were computed with the 2004 version of the Ten Tusscher–Noble–Noble–Panfilov (TNNP) membrane model of the human ventricular myocyte (3). The model included transmural and left-to-right ventricular heterogeneity of membrane properties. Computed activation times served as input for two simplified models:

- a "fixed-AP" model, which used predefined AP, and
- a "hybrid model," an RD model in which all cells were stimulated at fixed times, but AP shapes were computed.

Fixed APs were obtained from simulations of isolated cells, and represented the same heterogeneity of cell types as the heart model. The hybrid model was tested with spatial resolutions of 0.25, 0.5, and 1.0 mm. Electrocardiograms were computed with all three models.

Results

As expected, computed QRS complexes were practically identical in all models. Figure 1 shows that T waves in the fixed-AP model had 20 to 40% larger amplitudes in the precordial leads V1–V3, became biphasic in lead III, and notched in leads II and




Figure 1. Electrocardiogram simulated by the full reaction-diffusion model (black) and by the fixed-action potential model (red online/grey). Tick marks at the horizontal axis are placed at 100-ms intervals. The vertical scale is in milliVolts




Figure 2. Electrocardiogram simulated by the full reaction-diffusion model at 0.25-mm resolution (black) and by the hybrid model at 1-mm resolution (red online/grey). Tick marks at the horizontal axis are placed at 100-ms intervals. The vertical scale is in milliVolts

aVF. In contrast, the hybrid model produced the same T waves as the RD model at 0.25-mm resolution, but underestimated T wave amplitude in V1–V3 at lower resolutions. The case of 1-mm resolution is shown in Figure 2.

Discussion

Fixed AP waveforms in a forward ECG model can lead to exaggerated T waves. Hybrid models only equal RD models when used with the same spatial resolution. This presents only a very small efficiency improvement over a full RD model. Thus, accurate simulation of electrocardiographic T waves does require an RD model. In contrast, the QRS complex is not very sensitive to the shape of the AP, and can be simulated with any model that correctly represents activation times (6, 7). Only in case of severe propagation problems does this imply the use of an RD model.

The differences in T-wave shape that we observed are probably specific for strong transmural heterogeneity of repolarisation times. If such large gradients are not present in a model, the use of high spatial resolution is likely to be less important.

Our hybrid model used activation times determined by our RD model. Using exactly the same activation times allowed us to determine T-wave changes independent of activation order. In practice, of course, activation times for a hybrid model must be computed somehow. Typically, fixed propagation velocities are used for such purposes (6). Another interesting method is to use a modification of the membrane model that slows down the depolarizing sodium current to make the depolarization wavefront thicker and less steep (8). This allows the use of a lower spatial resolution. However, the side effects of such modifications are hard to predict, so this method too should be used with caution.

Acknowledgements

Computational resources for this work were provided by the Réseau québécois de calcul de haute performance (RQCHP). M. Potse was supported by a postdoctoral award from the Groupe de recherche en sciences et technologie biomédicale (GRSTB), École Polytechnique and Université de Montréal; and by the Research Center of Sacré-Coeur Hospital, Montréal, Québec, Canada.

References

1. Noble D, Rudy Y. Models of cardiac ventricular action potentials: iterative interaction between experiment and simulation. *Phil Trans Roy Soc London; Phys Sc* 2001; 359: 1127-42.
2. Iyer V, Mazhari R, Winslow RL. A computational model of the human left-ventricular epicardial myocyte. *Biophys J* 2004; 87: 1507-25.
3. ten Tusscher KHWJ, Noble D, Noble PJ, Panfilov AV. A model for human ventricular tissue. *Am J Physiol Heart Circ Physiol* 2004; 286: H1573–89.
4. Trudel MC, Dubé B, Potse M, Gulrajani RM, Leon LJ. Simulation of propagation in a membrane-based computer heart model with parallel processing. *IEEE Trans Biomed Eng* 2004; 51: 1319-29.
5. Potse M, Dubé B, Richer J, Vinet A, Gulrajani RM. A comparison of monodomain and bidomain reaction-diffusion models for action potential propagation in the human heart. *IEEE Trans Biomed Eng* 2006; 53: 2425-35.
6. Lorange M, Gulrajani RM. A computer heart model incorporating anisotropic propagation: I. Model construction and simulation of normal activation. *J. Electrocardiol* 1993; 26: 245-61.
7. Colli Franzone P, Guerri L. Spreading of excitation in 3-D models of the anisotropic cardiac tissue. I. validation of the eikonal model. *Math Biosci* 1993; 113: 145-209.
8. Bernus O, Verschelde H, Panfilov AV. Modified ionic models of cardiac tissue for efficient large scale computations. *Phys Med Biol* 2002; 47: 1947-59.